

Soil Quality Resource Concerns: **Soil Biodiversity**

USDA Natural Resources Conservation Service

January 1998

What is soil biodiversity?

Soil biodiversity reflects the mix of living organisms in the soil. These organisms interact with one another and with plants and small animals forming a web of biological activity.

Soil is by far the most biologically diverse part of Earth. The soil food web includes beetles, springtails, mites, worms, spiders, ants, nematodes, fungi, bacteria, and other organisms. These organisms improve the entry and storage of water, resistance to erosion, plant nutrition, and break down of organic matter. A wide variety of organisms provides checks and balances to the soil food web through population control, mobility, and survival from season to season.



What are the benefits of soil organisms?

Residue decomposition

Soil organisms decompose plant residue. Each organism in the soil plays an important role. The larger organisms in the soil shred dead leaves and stems. This stimulates cycling of nutrients. The larger soil fauna include earthworms, termites, pseudoscorpions, microspiders, centipedes, ants, beetles, mites, and springtails.

When mixing the soil, the large organisms bring material to smaller organisms. The large organisms also carry smaller organisms within their systems or as “hitchhikers” on their bodies.

Small organisms feed on the by-products of the larger organisms. Still smaller organisms feed on the products of these organisms. The cycle repeats itself several times with some of the larger organisms feeding on smaller organisms.

Some larger organisms have a life span of two or more years. Smaller organisms generally die more quickly, but they also multiply rapidly when conditions are favorable. The food web is therefore quick to respond when food sources are available and moisture and temperature conditions are good.

Infiltration and storage of water

Channels and aggregates formed by soil organisms improve the entry and storage of water. Organisms mix the porous and fluffy organic material with mineral matter as they move through the soil. This mixing action provides organic matter to non-burrowing fauna and creates pockets and pores for the movement and storage of water. Fungal hyphae bind soil particles together and slime from bacteria help hold clay particles together. The water-stable aggregates formed by these processes are more resistant to erosion than individual soil particles. The aggregates increase the amount of large pore space which increases the rate of water infiltration. This reduces runoff and water erosion and increases soil moisture for plant growth.

Nutrient cycling

Soil organisms play a key role in nutrient cycling. Fungi, often the most extensive living organisms in the soil, produce fungal hyphae. Hyphae frequently appear like fine white entangled thread in the soil. Some fungal hyphae (mycorrhizal fungi) help plants extract nutrients from the soil. They supply nutrients to the plant while obtaining carbon in exchange and thus extend the root system. Root exudates also provide food for fungi, bacteria, and nematodes.

When fungi and bacteria are eaten by various mites, nematodes, amoebas, flagellates, or ciliates, nitrogen is released to the soil as ammonium. Decomposition by soil organisms converts nitrogen from organic forms in decaying plant residues and organisms to inorganic forms which plants can use.

Management considerations

Cultivation

The effects of cultivation depend on the depth and frequency of the cultivation. Tilling to greater depths and more frequent cultivations have an increased negative impact on all soil organisms. No-till, ridge tillage, and strip tillage are the most compatible tillage systems that physically maintain soil organism habitat and biological diversity in crop production.

Compaction

Soil compaction reduces the larger pores and pathways, thus reducing the amount of suitable habitat for soil organisms. It also can move the soil toward anaerobic conditions, which change the types and distribution of soil organisms in the food web. Gaps in the food web induce nutrient deficiencies to plants and reduce root growth.

Pest control

Pesticides that kill insects also kill the organisms carried by them. If important organisms die, consider replacing them. Plant-damaging organisms usually increase when beneficial soil organisms decrease. Beneficial predator organisms serve to check and balance various pest species.

Herbicides and foliar insecticides applied at recommended rates have a small impact on soil organisms. Fungicides and fumigants have a much greater impact on soil organisms.

Fertility

Fertility and nutrient balances in the soil promote biological diversity. Typically, carbon is the limiting resource to biological activity. Plant residue, compost, and manure provide carbon. Compost also provides a mix of organisms, so the compost should be matched to the cropping system.


(Prepared by the National Soil Survey Center, NRCS, USDA in cooperation with the Soil Quality Institute, and the National Soil Tilth Laboratory, Agricultural Research Service, USDA).

The U. S. Department of Agriculture (USDA) prohibits discrimination in its programs on the basis of race, color, national origin, gender, religion, age, disability, political beliefs, sexual orientation, and marital or family status. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at (202) 720-2600 (voice and TDD).

To file a complaint of discrimination, write USDA, Director, Office of Civil Rights, Room 326W, Whitten Building, 14th and Independence Avenue, SW, Washington, DC 20250-9410 or call (202) 720-5964 (voice or TDD). USDA is an equal opportunity provider and employer.

Cover crops and crop rotations

The type of crops that are used as cover or in crop rotations can affect the mix of organisms that are in the soil. They can assist in the control of plant pests or serve as hosts to increase the number of pests. Different species and cultivars of crops may have different effects on pests. However, the organisms and their relation to the crop are presently not clearly understood.



Crop residue management

Mixing crop residue into the soil generally destroys fungal hyphae and favors the growth of bacteria. Since bacteria hold less carbon than fungi, mixing often releases a large amount of carbon as carbon dioxide (CO₂). The net result is loss of organic matter from the soil.

When crop residue is left on the soil surface, primary decomposition is by arthropod shredding and fungal decomposition. The hyphae of fungi can extend from below the soil surface to the surface litter and connect the nitrogen in the soil to the carbon at the surface. Fungi maintain a high C:N ratio and hold carbon in the soil. The net result is toward building the carbon and organic matter level of the soil. In cropping systems that return residue, macro-organisms are extremely important. Manage the soil to increase their diversity and numbers.

Visit our Web site:
<http://soils.usda.gov>